

Elachistocleis bicolor (Amphibia: Anura: Microhylidae) and *Eupemphix nattereri* (Amphibia: Anura: Leuiperidae): Morphological abnormalities from Reserva Natural Laguna Blanca, Paraguay

Jean-Paul Brouard¹, Paul Smith^{1,2}

¹ Para La Tierra, Reserva Natural Laguna Blanca, Santa Rosa del Aguaray, San Pedro, Paraguay. www.paralatierra.org

² Fauna Paraguay, Encarnación, Paraguay. www.faunaparaguay.com

Corresponding author. E-mail: jeanpaul@paralatierra.org

The global amphibian decline is well-documented, with the complex effects and interactions of ultraviolet radiation, predation, habitat modification, diseases, changes in climate or weather patterns, introduced predators, environmental acidity and toxicants all known to be causal factors (Alford and Richards, 1999). The drastic increase in the frequency of developmental abnormalities observed in recent years has also been linked to this phenomenon. Although the precise causes of such abnormalities are still imperfectly known, contamination from pollutants, viral and parasitic infections, ultraviolet radiation and predation in the early metamorphic stages, have all

been mooted as potential contributors (Lannoo, 2008). Morphological abnormality of 0-5% is commonly found in amphibian populations, but when a population exceeds this mark it is considered abnormally high (Bionda *et al.*, 2012; Piha *et al.*, 2010; Pekkonen and Merilä, 2006).

Reserva Natural Laguna Blanca (RNLB) is located in the Cerrado zone of Departamento San Pedro, northeastern Paraguay (Fig. 1). It is a small, 804 ha reserve based around a freshwater lake of 157 ha (Guyra Paraguay, 2008). The reserve comprises a tract of 400 ha of near pristine Cerrado, a degraded section of Atlantic forest and a transitional zone of semi-deciduous, semi-humid gallery forest (Eiten, 1972; Eiten, 1978).

Specimens of *Elachistocleis bicolor* (Guérin-Méneville, 1838) (Microhylidae) and *Eupemphix nattereri* (Steindachner, 1863) (Leuiperidae) with malformations were collected near a temporary pool (23°46'52.6"S, 56°17'28.9"W) in a zone of transition between Cerrado and Atlantic Forest habitats at RNLB. The specimens are housed in the Colección Zoológica de Para La Tierra (CZPLT), located at the reserve. Morphological abnormalities were identified using Meteyer's (2000) classification. Comparative morphometry of the normal limb vs. abnormal limb of the two *E. bicolor* specimens and the two abnormal forelimbs of the *E. nattereri* specimen are given in tables 1 and 2. These specimens account for the first documented records of morphological abnormalities in amphibians from Paraguay.

Elachistocleis bicolor is a fossorial microhylid with a wide distribution comprising southern Brazil, Bolivia, Paraguay, Uruguay and northern Argentina (Weiler *et al.*, 2013). It can be found in various eco-regions including Atlantic Forest, Cerrado, Pantanal, and humid and dry Chaco (Caramaschi, 2010; Brusquetti and Lavilla, 2006; Nuñez *et al.*, 2004). New species of *Elachistocleis* are currently being described with regularity (Caramaschi, 2010; Kwet and Di-Bernardo, 1998; Lavilla *et al.*, 2003; Nunes de Almeida and Toledo, 2012; Pereyra *et al.*, 2013; Toledo, 2010; Toledo *et al.*, 2010), but specimens ascribed to *E. bicolor* here were those with a uniform yellow ventral surface, no dorsomedial line, and a head length less than 90% of the head width, in accordance with Caramaschi (2010).

Three specimens of *Elachistocleis bicolor* show abnormalities of the limbs (Table 1). CZPLT-H-487 (3 June 2013; female; 32.56 mm snout-vent length [SVL]; Fig. 2A-B) and




Figure 1: Map showing location of Reserva Natural Laguna Blanca, Departamento San Pedro, Paraguay.




Figure 2: Morphological abnormalities described in the text from Reserva Natural Laguna Blanca, Departamento San Pedro, Paraguay. A-F: *Elachistocleis bicolor* specimens CZPLT-H-487 (A-B), CZPLT-H-521 (C-D) and CZPLT-H-584 (E-F). G-H: *Eupemphix nattereri* specimen CZPLT-H-586.

Table 1: Comparative morphometry of the normal limb vs. abnormal limb of *Elachistocleis bicolor* specimens. Measurements in mm.

	CZPLT-H-487 (abnormal)		CZPLT-H-521 (normal)	
	Left Side	Right Side	Left Side	Right Side
Femur	10.6	8.4	10.4	10.2
Tibiafibula	5.2	0	8.5	8.4
Foot	11.6	4.4	11.7	6.6

Table 2: Comparative morphometry of two abnormal forelimbs of the *Eupemphix nattereri* specimen 586. Measurements in mm.

	CZPLT-H-586	
	Left Side	Right Side
Humerus	12.0	12.1
Radio-ulna	3.1	10.6
Hand	0	4.2

CZPLT-H-521 (8 July 2013; male; 26.40 mm SVL; Fig. 2C-D) both have a reduced right hindlimb. CZPLT-H-487 displays hemimelia (reduction of the long bone) of the femur, an absent tibiafibula, ectrodactyly (absent digits) and brachydactyly (reduced digits) of the foot (three, reduced toes present compared to the norm of five), and rotation of the foot. CZPLT-H-521 shows ectrodactyly and brachydactyly. The femur, tibiafibula and tibiale/fibulare are present, but only two digits are complete. CZPLT-H-584 (29 November 2013; male; 23.04 mm SVL; Fig. 2E-F) has an abnormal right forelimb showing ectromelia of the tibiale/fibulare. The humerus and radio-ulna are complete, but the hand is reduced with no digits present.

Eupemphix nattereri is a widespread frog species distributed from central and southeast Brazil to eastern Paraguay and Bolivia (Weiler *et al.*, 2013). It can be found in a wide range of habitats including the humid Chaco, Cerrado, Atlantic Forest, and Pantanal (Brusquetti and Lavilla, 2006).

Specimen CZPLT-H-586 (29 November 2013; male; 47.7 mm SVL; Fig. 2G-H) has two abnormal forelimbs. The right limb shows ectromelia of the tibiale and fibulare (complete humerus and radio-ulna, but a reduced hand without digits), and the left limb shows ectromelia of the radio-ulna (complete humerus, reduced radio-ulna, and hand absent).

Despite ongoing year-round amphibian inventories at RNLB, morphological abnormalities have so far been discovered only on the limbs of these two species. Although the causes of the observed abnormalities are not immediately obvious, the low frequency of abnormalities and high level of species specificity, despite the large number of specimens and species collected, may be suggestive of a non-environmental cause, perhaps related to unsuccessful predation, accidental injuries or developmental errors (Guderyahn, 2006; Székely and Szilárd, 2003). All four specimens were adult and did not exhibit any obvious lack of fitness or detrimental health effects to suggest that such abnormalities might hinder their ability to survive.

ACKNOWLEDGMENTS

We thank the Secretaria del Ambiente (SEAM) for issuing the relevant permits and for their recognition of the importance

of RNLB by declaring it a Reserva Natural. Specimens were collected under permit 02/13. Particular thanks to the Duarte family, the owners of Laguna Blanca, for their foresight and support of PLT, without which the fieldwork would not have been possible. All volunteers and interns who contributed to the amphibian research by assisting in the field work deserve praise for their efforts.

REFERENCES

- Alford, R. A. and S. J. Richards. 1999. Global amphibian declines: A problem in applied ecology. *Annual Review of Ecology and Systematics*, 30: 133-165.
- Bionda, C., N. Salas, E. Caraffa, M. Baraquet and A. Martino. 2012. On abnormalities recorded in an urban population of *Rhinella arenarum* from central Argentina. *Herpetology Notes*, 5: 237-241.
- Brusquetti, F. and E. Lavilla. 2006. Lista comentada de los anfibios de Paraguay. *Cuadernos de Herpetología*, 20: 3-79.
- Caramaschi, U. 2010. Notes on the taxonomic status of *Elachistocleis ovalis* (Schneider, 1799) and description of five new species of *Elachistocleis* Parker, 1927 (Amphibia, Anura, Microhylidae). *Boletim do Museu Nacional, Rio de Janeiro, Zoologia*, 527: 1-32.
- Eiten, G. 1972. The Cerrado vegetation of Brazil. *Botanical Review*, 38: 201-341.
- Eiten, G. 1978. Delimitation of the Cerrado concept. *Vegetatio*, 36: 169-178.
- Guderyahn, L. 2006. Nationwide assessment of morphological abnormalities observed in amphibians collected from United States National Wildlife Refuges. U.S. Fish and Wildlife Service, Report Number CBFO-C06-01. 74pp.
- Guyra Paraguay. 2008. Áreas importantes para la conservación de las aves del Paraguay. Guyra Paraguay, Asunción, 470pp.
- Johnson, P. T. J. and J. Bowerman. 2010. Do predators cause frog deformities? The need for an eco-epidemiological approach. *Journal of Experimental Zoology (Molecular and Developmental Evolution)*, 314B: 515-518.
- Kwet, A. and M. Di-Bernardo. 1998. *Elachistocleis erythrogaster*, a new microhylid species from Rio Grande do Sul, Brazil. *Studies on Neotropical Fauna and Environment*, 33: 7-18.
- Lannoo, M. 2008. Malformed frogs: The collapse of aquatic ecosystems. University of California Press, Berkeley, USA. 270pp.
- Lavilla, E. O., M. Vaira and L. Ferrari. 2003. A new species of *Elachistocleis* (Anura: Microhylidae) From the Andean Yungas of Argentina, with comments on the *Elachistocleis ovalis* – *E. bicolor* controversy. *Amphibia-Reptilia*, 24: 269-284.
- Meteyer, C. U. 2000. Field guide to malformations of frogs and toads with radiographic Interpretations. Biological Science Report USGS/BRD/BSR-2000-0005. US Department of the Interior, Geological Survey. 18pp.
- Nunes-de-Almeida, C. H. L. and L. F. Toledo. 2012. A new species of *Elachistocleis* Parker (Anura, Microhylidae) from the state of Acre, northern Brazil. *Zootaxa*, 3424: 43-50.
- Núñez, D., R. Maneyro, J. Langone, J. and R. O. de Sá. 2004. Distribución geográfica de la fauna de anfibios del Uruguay. *Smithsonian Information Herpetological Service*, 134: 1-34.
- Pereyra, L. C., M. S. Akmentins, G. Laufer and M. Vaira. 2013. A new species of *Elachistocleis* (Anura: Microhylidae) from north-western Argentina. *Zootaxa*, 3694: 525-544.
- Piña, H., M. Pekkonen and J. Merilä. 2006. Morphological abnormalities in amphibians in agricultural habitats: A case study of the common frog *Rana temporaria*. *Copeia*, 2006(4): 810-817.
- Székely, P. and N. Szilárd. 2003. The incidence of mutilations and malformations in a population of *Pelobates fuscus*. *Russian Journal of Herpetology*, 10: 145-148.
- Toledo, L. F. 2010. A new species of *Elachistocleis* (Anura; Microhylidae) from the Brazilian Amazon. *Zootaxa*, 2496: 63-68.
- Toledo, L. F., D. Loebmann, C. F. B. Haddad. 2010. Revalidation and redescription of *Elachistocleis cesarii* (Miranda-Ribeiro, 1920) (Anura: Microhylidae). *Zootaxa*, 2418: 50-60.
- Weiler, A., K. Núñez, E. Lavilla, S. Peris and D. Baldo. 2013. Anfibios del Paraguay. FACEN, Asunción, Paraguay. 126pp.